



Floral abundance and resource quality influence pollinator choice

ROBERT E. FOWLER, ELLEN L. ROTHERAY* and DAVE GOULSON

School of Life Sciences, University of Sussex, Falmer, UK

Abstract. 1. Pollinator declines caused by forage habitat loss threaten insect pollination services. Pollinating insects depend on adequate floral resources, and their ability to track these resources. Variability of these resources and the effect on insect foraging choice is poorly understood.

2. We record patterns of visitation to six wildflower species and test the hypotheses that: pollinators preferentially visit the most rewarding flowers; nectar diurnal variations affect foraging preferences; pollinators respond most strongly to nectar rewards.

3. Nectar volume and sugar concentration were negatively correlated within plant species over time of day where greater concentration and lower volume was evident in the afternoon, but this did not correspond to pollinator visitation. Both floral abundance and nectar quality (total sugar per inflorescence) positively affect insect visitation. For some foragers, the positive effects of high-quality rewards were only evident when floral abundance was high (> 50 inflorescences per patch), perhaps reflecting the low probability of pollinators detecting scarce rewards. Pollen quality (total protein per inflorescence) was negatively related to visitation of *Apis mellifera* and *Bombus pascuorum*.

4. Fewer pollinators visiting flowers of higher pollen quality could reflect plant allocation trade-offs or the presence of secondary metabolites in pollen, meaning pollen foraging is likely affected by factors other than protein concentration. Nectar rather than pollen appeared to be the main driver of floral choice by insects in this system.

5. Conservation schemes for bees in farmland or gardens might benefit from ensuring that rewarding plant species are present at high density and/or are aggregated in space.

Key words. Bees, nectar, pollen, protein, sugar.

Introduction

Over the past 80 years, local and UK-wide changes in farming practice and agricultural intensification have led to a reduction in diversity of crop and non-crop plants (Robinson & Sutherland, 2002; Öckinger & Smith, 2007; Geiger *et al.*, 2010). This includes the loss of meadow plants (Goulson *et al.*, 2005), arable weeds and hedgerows (Hanley & Wilkins, 2015), which provide valuable foraging resources for flower visiting insects. This habitat

degradation has been identified as the primary reason for population declines in insects reliant on flowers to provide nectar and/or pollen as a main food source, including adult butterflies (Brereton *et al.*, 2011) and adult and offspring honey bees (Potts *et al.*, 2010), bumblebees (Goulson *et al.*, 2008) and solitary bees (Ollerton *et al.*, 2014). Nesting and hibernation resources aside, it is imperative that bees and other pollinating insects are able to forage effectively for nutritional resources in increasingly fragmented landscapes in order to survive and reproduce, particularly as they are facing other pressures such as diseases and pathogens (Cox-Foster *et al.*, 2007), global environmental change (Tylianakis *et al.*, 2008) and pesticide use (Goulson *et al.*, 2015), all of which impact on their survival. Population decline in flower visiting insects

Correspondence: Robert E. Fowler, School of Life Sciences, University of Sussex, Falmer BN1 9QG, UK.
E-mail: R.E.Fowler@sussex.ac.uk

*Co-first author.

could jeopardise the pollination services provided to entomophilous crops (Kremen *et al.*, 2002; Klein *et al.*, 2007).

Nectar provides sugars (mainly sucrose, glucose and fructose), which energise pollinators to continue foraging (on nectar or other sources of nutrients), undertake nesting activities, find mates, and provide for offspring. It also contains ions, water and small amounts of amino acids, which may contribute to nutrition (Kim & Smith, 2000). Pollen comprises proteins, carbohydrates, lipids, vitamins and minerals (Roulston & Cane, 2000; Nicolson, 2011). Although many flower visiting insects consume pollen for sustenance (e.g. beetles, flies), bees (Hymenoptera: Apoidea) also collect pollen to feed larvae, and many simultaneously collect both nectar and pollen from flowers, depending on what requirements they have and what flower species they are feeding on (Heinrich, 1979a,b; Goulson *et al.*, 2005).

Survival and reproductive success of pollinating insects is dependent on them successfully gathering adequate protein and sugar to provide for their energy and nutritional needs. For example, many Lepidoptera, Diptera and Hymenoptera rely on energy gained from nectar to undertake mating flights, for dispersal and/or migration, and to find suitable places to lay their eggs. Butterflies tend to exhibit lower fecundity when nectar is limited (Boggs & Ross, 1993), and the hoverfly *Episyrphus balteatus* has greater longevity when fed a sugar and protein-rich diet (Pinheiro *et al.*, 2015). Bees require both pollen and nectar to feed to their offspring. When fed on protein-rich diets, bumblebee colonies are more reproductively successful (Génissel *et al.*, 2002; Kitaoka & Nieh, 2009), and the solitary bee *Lasioglossum zephyrum* (Roulston & Cane, 2002) and honey bees (Basualdo *et al.*, 2012) produce larger adults. Larger bees generate and retain heat faster leading to earlier and more frequent forage flights (Stone, 1993), they have greater diapause survival (Strohm & Linsenmair, 1997) and are better able to cope with adverse conditions such as parasitism and disease (Alaux *et al.*, 2010; Di Pasquale *et al.*, 2013).

In order to forage effectively, pollinators use olfactory cues to enable detection of non-depleted nectar resources (Goulson *et al.*, 1998a; Howell & Alarcón, 2007) and greater nectar volume and sugar content (Pyke, 1978; Heinrich, 1979a; Wolff, 2006). Likewise, insect foragers tend to select pollen with greater protein and essential amino acid content (Levin & Bohart, 1955; Schmidt, 1982; Robertson *et al.*, 1999; Cook *et al.*, 2003; Arenas & Farina, 2012), and obligate insect pollinated plant species produce pollen that is richer in protein and amino acids compared to facultative species (Hanley *et al.*, 2008; but see Roulston *et al.*, 2000).

Few studies have investigated how eusocial or solitary bees integrate information on nectar and pollen quantity and quality simultaneously, and those that have, indicate that nectar is the primary factor influencing foraging preference for bees, and pollen a secondary consideration (Konzmann & Lunau, 2014; but see Somme *et al.*, 2015). In addition, studies focusing on flower selection by

pollinators tend to look only at individual species of bee and specific pollinators for individual plant species (e.g. Robertson *et al.*, 1999), or solely bees as a group (e.g. Heinrich, 1979b; Pernal & Currie, 2001). Floral choice tests are frequently undertaken in controlled conditions (e.g. Konzmann & Lunau, 2014), but the limited number of field studies investigating the influence of rewards on the foraging decision of pollinators makes it difficult for the conclusions gained by laboratory studies to be applied, especially as there are external factors in the natural environment likely to influence the results, such as the spatial distribution and reward phenology of flowers. For example, Hanley and Wilkins (2015) describe greater food plant abundance in roadside, compared with field facing hedgerows, and noted a corresponding increase in bumblebee abundance. In addition, several studies indicate that some flower species offer less nectar in the middle of the day and afternoon compared to the morning and evening (Mačukanović-Jocić *et al.*, 2004; Silva & Dean, 2004; Mačukanović-Jocić & Đurđević, 2005), providing a further challenge to efficient foraging by pollinators since the relative rewards provided by different flower species may alter hour by hour through the day.

In this study, we examine the foraging choices made by all flower visiting insects in relation to the relative nectar and pollen quantity and quality of six common plant species under natural conditions. We tested the hypothesis that nectar quantity or quality shows diurnal variation and that nectar or pollen quantity or quality, or a combination of reward metrics, predicts insect visits. We record how nectar volume and sugar concentration changes through the day, and also the pollen weight and protein concentration for each test plant species, and relate these values to insect flower choices through the day. Specifically, this study's objectives were to; (i) record nectar and pollen quantity and quality estimates for six test plant species, (ii) investigate the diurnal variation in nectar quantity and quality for the test plant species, and (iii) assess how nectar and pollen quantity and quality influence foraging choices by flower visiting insects.

Methods

Test plant species

Six species of flowering plant that are common in southern UK were selected for nectar and pollen quantity and quality estimation, and insect visitation surveys in the field. These were *Lamium album* L. and *Glechoma hederacea* L. (Lamiaceae), *Crataegus monogyna* Jacq. and *Rubus fruticosus* L. agg. (Rosaceae), *Symphytum officinale* L. (Boraginaceae) and *Ranunculus repens* L. (Ranunculaceae). These spring or early summer flowering plants considered to be beneficial foraging plants for insects in previous studies (Lack, 1982; Goulson *et al.*, 1998b; Kipling & Warren, 2013) (further species details are listed in Table S1 in the Electronic Supplementary Material).

Previously, *R. fruticosus* and *S. officinale* were included in studies of pollen protein quality and its effects on bee foraging preferences (Hanley *et al.*, 2008), but to the best of our knowledge, the other test plant species have not been studied in this context.

Site selection

Sites were chosen to be included in this study if they contained at least three of the test plant species within 50 m of each other, were on chalk soil, were easy to access and were subject to intermediate levels of disturbance (e.g. mowed once a year, or grazed intermittently by sheep). Sites could be divided into two groups, road verges and semi-improved grassland. In terms of management, road verge sites were mowed once a year and semi-improved grassland were grazed several times throughout the year. None of the survey sites were either mown or grazed while this study was taking place. Once sites were identified, nectar and pollen samples were collected from test plant species and visitation sampling was conducted.

Nectar volume and sugar concentration

Nectar sampling was undertaken on 30 inflorescences of each test plant species, collected from at least three different survey sites. Nectar was sampled from inflorescences of plant species that had recently come in to bloom and were easily accessible. For each inflorescence, nectar production was estimated from morning, afternoon and evening periods for each plant species tested. For each species, then, flowers were emptied of nectar and bagged by 09:00 (GMT) using a fine-mesh cotton fabric and masking tape. At 15:00, 21:00 and 09:00, the following morning flowers were emptied of nectar using 5 μ l micropipettes (BLAUBRAND intraMARK, Wertheim, Germany). The nectar from each inflorescence was then expelled onto a field refractometer (Bellingham and Stanley Ltd, Basingstoke, UK) to measure the proportion of sugar in each sample (Bolten *et al.*, 1979). This produced both mean nectar volume and mean sugar concentration for each species in each time period, and over a 24 h period when totalled.

Pollen weight and protein concentration

Pollen was collected from 10 inflorescences from at least three different sites for each plant species. Standing crop of pollen was taken just once for each replicate so variation in pollen quantity or quality was not measured over time of day. Flowers were collected in the field before anthers had dehisced, and placed in water in an unheated and well-ventilated laboratory until anthers opened. Pollen was collected systematically until anthers were empty. It is conceivable that placing flowers in water affects

pollen quality, but all plant species were treated in the same way. Pollen was either stored in a -20°C freezer for drying at a later date, or immediately dried in an oven for 24 h at 40°C . After drying, pollen was weighed to measure total pollen weight per inflorescence. Protein extraction and detection was undertaken using the Bradford assay (Bradford, 1976). The assay binds protein to Coomassie Brilliant Blue G-250 dye (4.7% [weight: volume] ethanol, and 8.5% [weight: volume] phosphoric acid dissolved in water). Light absorbance is then measured against known protein standards. From each of the 10 inflorescence pollen from each species, 1 mg pollen were dusted with aluminium powder, wetted with 20 μ l 0.1 mol L⁻¹ NaOH and ground with a micropestle. Ground pollen was reanimated with 480 μ l 0.1 mol L⁻¹ NaOH and placed in a refrigerator for 24 h before analysis, but used within 1 week. Prior to absorbance, measurement samples were placed in boiling water for 5 minutes and centrifuged for 5 minutes. Then, 10 μ l of supernatant was slowly vortexed with 300 μ l of dye reagent. This was repeated in triplicate for each sample and left to incubate at room temperature for 15 minutes.

Protein standards were made up each time samples were run, using pre-mixed concentrations of Bovine Serum Albumin (BSA) from the BIO-RAD (Hertfordshire, UK) Quick-Start Bradford Protein Assay kit. Once samples and standards were created, they were measured for absorbance within an hour of mixing at 595 nm using a Thermo Scientific (Paisley, UK) Nanodrop 2000 UV-Vis Spectrophotometer. This produced the standing crop of both mean pollen weight and a crude mean protein concentration for each test plant species.

Visitation surveys

Visitation surveys took place at seven sites across Sussex, UK between May and June 2014 (for site details see Table S2). Surveys were undertaken in the morning (08:00–10:00), afternoon (13:00–15:00) and evening (19:00–21:00). Each survey consisted of a 10 minute standing observation of a 4 m² area of each test plant species at each site. The number and species of all visiting insects to that plant species were recorded. All observed flower visiting insects appeared to collect nectar and, in most cases, pollen from plant species, therefore we did not attempt to distinguish which resources insects were collecting. The number of inflorescences of the plant species, ambient temperature and wind speed were recorded for each survey. Surveys were only conducted when air temperature was $>14^{\circ}\text{C}$ and average wind speeds was <20 mph.

Common species of bumblebee (genus *Bombus*) and European honeybees (*Apis mellifera* L.) were identified on the wing. Due to the unreliability of morphological characters for separating *Bombus terrestris* L. and *B. lucorum* agg. workers in the field, these two species were grouped. Although others have recorded flower visiting bumblebees

into groups based on colour type (Haughton *et al.*, 2003), in this study, the bumblebee species observed were easily separated to species in the field. Faded or suspected cleptoparasitic bees (*Psithyrus* spp.) were caught and checked using a hand magnifying glass. Other visiting insects too small to be identified on the wing were collected and identified to species or genus. Although unlikely given the short period of each survey, the possibility of double counting insects was the same for all surveys due to equal sampling time and therefore assumed to be unbiased across the study.

Statistical analysis

Means tests of variance were used to assess differences in floral resources available from the plant species, as well as the number of insect visits, between periods of the day. Based on whether resource and insect visit metrics showed homogeneity of variance between species or periods of the day using Levene's test, either ANOVA (parametric) or Kruskal–Wallis (non-parametric) tests were applied to assess variance for all groups before either Tukey HSD tests or pairwise Wilcox tests were applied to assess differences between groups respectively.

Generalised Linear Mixed Models (GLMMs) were used to test the effects of resource metrics on insect visitation (i.e. counts per 10 minute survey). Prior to applying models, proposed explanatory variables were checked for multicollinearity using Variance Inflation Factors (VIFs), if variables had VIFs >3 or correlation coefficients more than 0.6 with other variables, they were excluded from models (Zuur *et al.*, 2010). These analyses indicated that nectar volume and pollen weight were correlated with sugar and protein concentration respectively. To enable modelling of resource quality and quantity without violating model assumptions, nectar volume and pollen weight were multiplied by sugar and protein concentration to create total sugar and protein in milligrams per inflorescence as measurements of quality in the plant species. To account for the change in density of the solution based on the amount of sugar recorded, percentage sugar was multiplied by the mass density (g cm^{-3}) of sugar at that concentration. These measurements also allowed the interpretation of the nutritional gain available to foraging insect pollinators, and are referred to as total sugar and total protein from here on.

GLMMs were initially run with the environmental variables temperature and wind speed included as explanatory variables, however, these had no effect on visitation or the outcome of the models, so were removed. We modelled five visitation response variables (including: all insects; all bumblebees; the three most recorded species, *B. pascuorum* Scopoli, *B. pratorum* L., and *A. mellifera*) in response to the number of inflorescences (log transformed) for each test plant species surveyed in each observation area (4 m^2) (floral abundance), and total sugar and protein. We did not analyse data on other groups of

flower visiting insects alone as they were recorded in such small numbers that data analysis would have been unreliable. They were included in the data analysis as part of 'all insects' in statistical models, however. We included interactions between floral abundance and total sugar and protein explanatory variables. Sample site was fitted as a random factor because observations were nested within sample sites and contained different combinations of test plant species. Visitation rates were count data, therefore, models were applied with Poisson errors (O'Hara & Kotze, 2010). Model residuals were assessed for normality and heteroscedasticity. Model simplification was carried out using likelihood ratio tests, and sequentially deleting terms that did not significantly decrease model deviance, beginning with higher order interactions.

To further investigate how particular plant species affect insect visitation, linear mixed effects (LME) models were used to test the effects of floral abundance and species on the same response variables as the GLMMs, with site as a random factor. Models were tested for significance using likelihood ratio tests, with and without species as an explanatory variable.

Already published protein estimates for each plant species were compared to confirm the extraction results and accuracy of our values. We found that only *Lamium album* (4.48%) differed from the literature for *Lamium* sp. (22.8% in Roulston *et al.*, 2000), so to assess whether this estimate affected our findings, we substituted it in the two models that included total protein (for *A. mellifera* and *B. pascuorum* visitation).

Statistical analysis was conducted using R v3.1.2 (R Core Team, 2014) within RStudio (RStudio, 2012) using packages 'lme4' (Bates *et al.*, 2015) and 'usdm' (Naimi, 2013), and plots were generated using 'ggplot2' (Wickham, 2009).

Results

Nectar and pollen quantity and quality

A large amount of between-inflorescence variation was found in nectar and pollen resources for each test plant species (Table 1). Nevertheless, nectar volume (Kruskal–Wallis $H = 211.64$, d.f. = 5, $P < 0.001$), nectar sugar concentration ($H = 112.61$, d.f. = 5, $P < 0.001$) and pollen protein concentration (ANOVA $F_{5,54} = 20.6$, $P < 0.001$) varied significantly between plant species, with only pollen weight ($F_{5,54} = 1.25$, $P = 0.3$) showing no significant differences.

Over 24 h, *Rubus fruticosus* produced the greatest volume of nectar, which contained the lowest sugar concentration (Table 1). In contrast, *Ranunculus repens* produced the least nectar, while *Crataegus monogyna* offered the highest sugar concentration. *Lamium album* pollen offered the greatest mass per inflorescence, but had the lowest protein concentration of all species. *Crataegus monogyna* produced the least pollen mass while *G. hederacea* offered

Table 1. Measurements per inflorescence of nectar (30 replicates) and pollen (10 replicates) quality and quantity (Mean \pm SD) for *Crataegus monogyna*, *Rubus fruticosus*, *Glechoma hederacea*, *Lamium album*, *Ranunculus repens* and *Symphytum officinale* between times and totalled over 24 h from seven sites in Sussex, UK.

Test plant species	Nectar (μ l)			Nectar (%)			Sugar per Infl. (mg)			Protein per Infl. (mg)							
	Morning	Afternoon	Evening	Over 24 h	Morning	Afternoon	Evening	Morning	Afternoon		Evening	Over 24 h					
													Pollen (mg)	Pollen (%)	Over 24 h		
<i>Crataegus monogyna</i>	1.1 \pm 1.7	0.4 \pm 0.4	0.4 \pm 0.5	1.9 \pm 2.0	26.13 \pm 2.26	54.95 \pm 3.15	36.54 \pm 1.81	35.03 \pm 9.42	0.11 \pm 0.03	15.39 \pm 2.24	0.3 \pm 0.2	0.6 \pm 0.1	0.6 \pm 0.9	0.6 \pm 0.1	0.3 \pm 0.2	1.2 \pm 1.2	0.02 \pm 0.01
<i>Rubus fruticosus</i>	2.9 \pm 2.8	0.9 \pm 1.2	3.1 \pm 2.1	6.8 \pm 3.7	9.48 \pm 0.52	24.41 \pm 1.87	18.16 \pm 1.26	17.13 \pm 4.72	0.48 \pm 0.23	12.6 \pm 2.28	0.6 \pm 0.5	0.3 \pm 0.3	0.3 \pm 0.2	0.3 \pm 0.3	0.6 \pm 0.5	1.3 \pm 0.7	0.06 \pm 0.02
<i>Glechoma hederacea</i>	1.2 \pm 0.6	0.7 \pm 0.4	0.8 \pm 0.4	2.6 \pm 0.8	29.03 \pm 1.20	30.40 \pm 1.57	29.43 \pm 1.43	29.86 \pm 4.56	0.89 \pm 0.29	19.07 \pm 3.46	0.2 \pm 0.1	0.2 \pm 0.1	0.4 \pm 0.2	0.2 \pm 0.1	0.2 \pm 0.1	0.9 \pm 0.2	0.17 \pm 0.06
<i>Lamium album</i>	1.1 \pm 0.4	0.3 \pm 0.2	0.8 \pm 0.5	2.2 \pm 0.6	19.76 \pm 0.98	32.88 \pm 2.13	27.67 \pm 1.38	25.88 \pm 7.61	1.22 \pm 0.33	4.48 \pm 1.28*	0.2 \pm 0.1	0.2 \pm 0.1	0.2 \pm 0.1	0.2 \pm 0.1	0.3 \pm 0.2	0.6 \pm 0.3	0.06 \pm 0.02
<i>Ranunculus repens</i>	0.1 \pm 0.1	0.1 \pm 0.1	0.0 \pm 0.0	0.2 \pm 0.2	22.79 \pm 2.25	28.00 \pm NA	29.5 \pm 2.60	24.50 \pm 8.13	0.76 \pm 0.3	12.06 \pm 1.50	0.0 \pm 0.0	0.1 \pm NA	0.0 \pm 0.0	0.1 \pm NA	0.0 \pm 0.0	0.1 \pm 0.0	0.09 \pm 0.04
<i>Symphytum officinale</i>	1.8 \pm 0.8	1.3 \pm 0.5	1.2 \pm 0.6	4.3 \pm 1.1	24.07 \pm 1.06	44.59 \pm 0.77	33.25 \pm 1.61	33.88 \pm 4.60	0.95 \pm 0.31	9.71 \pm 1.38	0.5 \pm 0.2	0.7 \pm 0.2	0.5 \pm 0.2	0.7 \pm 0.2	0.5 \pm 0.2	1.7 \pm 0.4	0.09 \pm 0.03

*Bold values indicate the total for nectar measurements over 24 h, and the total for pollen measurements (for which there was only a single measure per replicate). This value differed from the previously recorded value of 22.8% in Roulston *et al.* (2000).

the highest protein concentration. We found no correlation between nectar and pollen quantity (Pearson $R = -0.46$, $P = 0.26$) or quality (Pearson $R = -0.33$, $P = 0.52$).

Nectar volume and pollen weight were multiplied by sugar and protein concentration, respectively, to create total sugar and protein in milligrams per inflorescence (Table 1). Over 24 h, *Symphytum officinale* produced the greatest total sugar per inflorescence while *R. repens* produced the least. *Crataegus monogyna* contained the most total protein per inflorescence and *S. officinale* the least (Table 1).

Diurnal variation in nectar quantity and quality

Mean nectar volume and sugar concentration averaged between plant species (regardless of time period) were not significantly correlated (Pearson $R = -0.49$, $P = 0.32$, Fig. 1a). When mean for each species at each time period was taken, there was a significant negative correlation between nectar volume and sugar concentration (Pearson $R = -0.52$, $P = 0.02$, Fig. 1b).

Volumes of nectar produced differed markedly between species, and between morning, afternoon and evening periods. *Rubus fruticosus* produced the greatest volume of nectar of all species in the evening and morning, yet recorded the lowest sugar concentration of all species in these periods (Table 1). The lowest volume of nectar was detected in *R. repens* in the evening, and the greatest sugar concentration was found in *C. monogyna* in the afternoon. When averaged across species, mean nectar volume did not differ significantly between periods of the day ($F_{2,15} = 1.18$, $P = 0.35$). Mean nectar sugar concentration across species increased significantly in the afternoon compared to the morning ($F_{2,15} = 4.01$, $P = 0.04$) (Fig. 2).

Once sugar and protein concentrations were calculated between time periods, total sugar was significantly lower in the afternoon period compared to the morning or evening for *R. fruticosus*, *G. hederacea*, *L. album* and *R. repens* (Fig. 3). The exceptions were *C. monogyna* which produced comparable amounts of total sugar in each time period sampled, and *S. officinale* which had marginally more total sugar in the afternoon compared to morning and evening periods.

Foraging choices of flower visiting insects

In total, we made 112 ten-minute observations (38, 39 and 35 in morning, afternoon and evening periods respectively), between May and June 2014 at seven sites in Sussex, during which 574 insects were recorded visiting test plant species. Proportionately, of all insect visits, 93% were by bees (Hymenoptera: Apidae), 6% by Diptera (<1% of which were hoverflies [Syrphidae]), <1% by Lepidoptera and <1% by Vespidae (Table S3). Of all flower visiting insects, the three most numerous were the bee




Fig. 1. Negative correlation between nectar volume (μl) and nectar sugar concentration (%) averaged between (a) test plant species (Pearson $R = -0.49$, $P = 0.32$) and (b) test plant species and time period (Pearson $R = -0.52$, $P = 0.02$). ▼, *Symphytum officinale*; ▲, *Crataegus monogyna*; ■, *Glechoma hederacea*; ◆, *Lamium album*; ●, *Rubus fruticosus*; +, *Ranunculus repens*. Black symbols = Morning (09:00 GMT), dark grey symbols = Afternoon (15:00) and light grey symbols = Evening (21:00) periods of nectar sampling.

species *Bombus pascuorum*, *B. pratorum*, and *Apis mellifera*, which represented 33%, 30% and 16% of all insect visits respectively (Table S3). The majority of insects were recorded visiting *R. fruticosus* (47%), *S. officinale* (24%) and *L. album* (19%) (Table S3). LME models showed significant species-specific preferences for the most commonly recorded flower visiting insects (Table S4). *B. pascuorum* mainly visited *L. album* and *S. officinale* (49% and 31% of this species' visits respectively), *B. pratorum* preferred *R. fruticosus* and *S. officinale* (57% and 36%), and *A. mellifera* mainly visited *R. fruticosus* (87%) (Fig. S1; Table S3).

Fewer insects were recorded in the evening (122) than the morning (221) and afternoon periods (231), and

differences in abundance were also evident within and between recorded insect and test plant species (Table S3). These differences were not significant when tested between surveys for all insect abundance ($F_{2,109} = 2.59$, $P = 0.07$). No significant difference was found between any response variable between different periods of the day when all test plant species were grouped or tested separately, except *G. hederacea* which had significantly more insects visiting during the afternoon compared to morning and evening periods ($F_{2,24} = 9.03$, $P = 0.001$; Fig. 4). This however does not correspond with the timing of maximum sugar availability, which appeared highest in the morning or evening for most species (Fig. 3). As standing crop of pollen was taken just once for each replicate, we could not quantify variation in pollen quantity or quality over the day.

Overall, flower abundance had a positive effect on the total number of bumblebee (bees within the genera *Bombus*) and all insect visitation (Table 2). Total sugar predicted insect visitation, with this relationship being positive for all response categories apart from *B. pascuorum* which was negative (Table 2). There were also a number of significant higher order interactions between floral abundance and total sugar and protein. For *Apis mellifera*, all bumblebees and all insects, the relationship between insect visitation and total sugar was weak or absent at low floral abundance (10–50 inflorescences per 4 m^2) but positive at high floral abundance (51–1000 inflorescences per 4 m^2) (Fig. 5). Total protein had a significant negative effect on *A. mellifera* visitation (Table 2). In addition, a negative interaction between total protein and floral abundance was found for *B. pascuorum* (Table 2), with the relationship weak or absent at low floral abundance (10–50 inflorescences per 4 m^2) but negative at high floral abundance (51–1000 inflorescences per 4 m^2). The three plant species most visited by insects (*R. fruticosus*, *S. officinale* and *L. album*) were also recorded as producing the lowest total pollen protein, however. When the protein value for *L. album* was replaced with the higher value reported in Roulston *et al.* (2000), there was no longer a negative significant effect of total protein on *B. pascuorum* visitation, and we found a positive significant effect of total sugar (Table S5).

Discussion

Many insects forage on flowering plants to gain key nutritional resources, largely nectar and pollen. What factors determine when plants secrete nectar is still largely unknown, as both internal and external factors can affect the rate of secretion (Heil, 2011). In our study, floral rewards of nectar and pollen (apart from pollen weight) significantly differed between test plant species. Similarly, nectar resources varied between periods of the day, while the single measure of pollen for each replicate meant this resource could not be tested for diurnal differences. Nectar volume and sugar concentration were negatively




Fig. 2. The mean nectar volume (μl) and mean nectar sugar concentration (%) measured from 30 inflorescences in Morning (09:00 GMT), Afternoon (15:00) and Evening (21:00) time periods averaged from six test plant species. Significant differences between time periods were identified using post hoc tests; time periods that do not share a letter show significant variation ($P < 0.05$), ns, no significance.




Fig. 3. Total sugar per inflorescence (mg) of test plant species at Morning (09:00 GMT), Afternoon (15:00) and Evening (21:00) time periods, with the test statistic (' F ' for ANOVA and ' H ' for Kruskal–Wallis respectively) and significance levels for each analysis of variance between periods of the day for each species. Significant differences between time periods were identified using post hoc tests; time periods that do not share a letter show significant variation ($P < 0.05$), ns, no significance.




Fig. 4. Mean insect visits to test plant species in Morning (08:00–10:00 GMT), Afternoon (13:00–15:00) and Evening (19:00–21:00) time periods, with the test statistic and significance levels of ANOVAs between periods of the day for each species. Significant differences between time periods were identified using post hoc tests; time periods that do not share a letter showed significant variation ($P < 0.05$).

Table 2. Generalised Linear Mixed Models (GLMM) test results investigating the effects of floral abundance, total nectar and pollen quality and interactions between floral abundance and each quality metric on visitation of insect foragers from seven sites in Sussex, UK.

Visitation	Explanatory variables left in model	Parameter estimate \pm SE	z-value	P-value	AIC	logLik
<i>Apis mellifera</i>	Log flower abundance	-0.066 ± 0.253	-0.261	0.794	331.2	-159.6
	Total sugar	7.845 ± 3.288	2.386	0.017		
	Total protein	-2.491 ± 0.927	-2.687	0.007		
	Log flower abundance \times Total sugar	-1.405 ± 0.639	-2.201	0.028		
<i>Bombus pratorum</i>	Total sugar	2.838 ± 0.424	6.687	< 0.001	423.2	-208.6
<i>Bombus pascuorum</i>	Log flower abundance	0.281 ± 0.249	1.125	0.260	419.5	-203.7
	Total sugar	-1.380 ± 0.453	-3.048	0.002		
	Total protein	-12.584 ± 4.158	-2.785	0.005		
	Log flower abundance \times Total protein	2.879 ± 0.888	3.241	0.001		
All <i>Bombus</i> abundance	Log flower abundance	0.849 ± 0.168	5.049	< 0.001	605.5	-297.7
	Total sugar	6.275 ± 1.889	3.322	< 0.001		
	Log flower abundance \times Total sugar	-1.074 ± 0.366	-2.932	0.003		
All insect abundance	Log flower abundance	0.425 ± 0.124	3.404	< 0.001	705.7	-347.9
	Total sugar	4.741 ± 1.522	3.113	0.001		
	Log flower abundance \times Total sugar	-0.776 ± 0.296	-2.623	0.008		

Displayed are parameter estimates \pm standard error, z-value and P-value for each explanatory variable in the final model after simplification, and model AIC and log-likelihood tests. P-values are in bold if < 0.05 significance.




Fig. 5. Mean (\pm standard deviation) visits per test plant species to illustrate the positive effect of greater total sugar on visitation in *Bombus* spp. (above), and total sugar on *Bombus* spp. visits (below) when separated between sample plots with high or low floral abundance; low floral abundance (10–50 inflorescences per a 4 m², left) shows little or no trend, while high floral abundance (51–1000 inflorescences, right) shows a positive trend. Grey lines and polygons indicate model best fit and 95% Confidence Interval.

correlated in test plant species, with sugar concentration greater in the afternoon, which corresponds with previous studies showing similar trends (Mačukanović-Jocić *et al.*, 2004; Silva & Dean, 2004; Mačukanović-Jocić & Đurđević, 2005). This may be due to decreased water availability or flowers reabsorbing nectar in the afternoon

(Silva & Dean, 2004) when humidity decreases and temperatures rise (Silva & Dean, 2004; Mačukanović-Jocić & Đurđević, 2005). While our results suggest that nectar resources vary in quantity and quality across the day, insect visitation did not track nectar availability (with the exception of *G. hederacea* where the opposite was found:

more insects were recorded visiting this species in the afternoon when lower total sugar was recorded). This may be an indication that some pollinators are not capable of accurately responding to changes in nectar production throughout the day, or it may be that their activity is constrained by other factors such as temperature. It is important to note that our methods meant that afternoon and evening nectar sampling occurred at 6-h intervals, compared with morning sampling which was undertaken 12 h after the flowers had been emptied and bagged the previous evening. We considered this as providing a fair estimate of how much nectar is likely available to foraging insects by the morning, however, the effect of early morning foraging between 05:00 and 09:00 GMT was not represented by our data and this should be taken into account in interpretations.

Due to the high energy cost of foraging, successfully selecting the most rewarding flowers is predicted to have a large impact on survival and reproductive success, especially when floral resources are fragmented. The most documented resource offered to insects by plants as an attractant for pollination is nectar. Foraging insects are capable of learning nectar rewards gained from visited flowers (Pyke, 1978), preferring to forage on flowers with significantly more nectar (Heinrich, 1979a; Wolff, 2006) and with greater sugar concentration (Hendriksma *et al.*, 2014). After testing for relationships between nectar and pollen resources and pollinator visitations, our results support this, as typically, we found that greater nectar resources had a positive effect on insect and bee visitation. Single species specialism in insect-plant mutualisms is rare (Waser *et al.*, 1996), and the majority of flower visiting insects have flexible foraging choices. Foraging bumblebees show flower constancy ('majoring' on one particular species of known reward) and flower infidelity ('minoring' on other flowers checking reward change over time) (Heinrich, 1979b). This behaviour allows foragers to track resources in multiple flower species in a habitat. In our study, the positive effects of high floral rewards, i.e. sugar, were often only apparent when the floral abundance of test plant species was high (>50 inflorescences), with scarce flowers tending to be visited less frequently even when comparatively highly rewarding. Our results appear to support Heinrich (1979b); if a flower species is both abundant and rewarding, then insects are very likely to have discovered its value and preferentially visit it. In addition, where there are more flowers in a patch, there is a greater total quantity of nectar. Hence, although quality has an influence on forager choice, the abundance of floral rewards in the local environment is important in insect selection of resources.

There is evidence that bees show preferences for pollen with higher protein (Levin & Bohart, 1955; Robertson *et al.*, 1999) and essential amino acid content (Cook *et al.*, 2003), and these preferences are supported by obligate insect pollinated plant species producing pollen that is richer in protein and amino acids (Hanley *et al.*, 2008). Other studies describe contrasting results where protein-

rich pollen seems to be preferred in some cases, but not in others (Wille *et al.*, 1985 in Praz *et al.*, 2008; Roulston & Cane, 2002). Our results suggest greater total protein was negatively related to visitation by *A. mellifera*, and for *B. pascuorum*, the negative relationship between protein content and visitation was more apparent when the floral abundance of test plant species was high (>50 inflorescences). There are several possible explanations. Firstly, although not significant, there was a negative relationship between nectar and pollen quantity per inflorescence (correlation coefficient -0.46), so if bees are basing decisions primarily on nectar rewards, they will tend to visit flowers with less pollen. We did not attempt to discern whether bees were collecting pollen only, nectar only, or both, but previous studies suggest that the majority of visits are for nectar (e.g. Goulson *et al.*, 2005). Second, bees may have been responding to other nutritional compounds present in pollen. Plant species may be making trade-offs between protein and other nutritional elements that drive foraging preferences such as sterols (Somme *et al.*, 2015), lipids and starch (Roulston & Cane, 2000) or pollen-specific odours (Dötterl & Vereecken, 2010) not addressed in this study. Third, some plant species protect their pollen with defensive secondary compounds that may affect bee foraging choices (Gosselin *et al.*, 2013). For example, *Echium vulgare* has high protein content and also high concentrations of the hepatotoxins 1,2-dehydropyrrolizidine alkaloids and their N-oxides (Boppré *et al.*, 2005), which can be toxic to insects (e.g. Boppré *et al.*, 2005; Sedivy *et al.*, 2012) and affect flower selection (Kessler & Halitschke, 2009). In this study, the species recorded with the greatest amount of pollen protein were *Glechoma hederacea* and *Ranunculus repens*, and pollen of *Ranunculus* spp. is known to contain the toxic lactone protoanemonin ranunculin with reportedly negative effects on honeybees (Jurgens & Dötterl, 2004; Sedivy *et al.*, 2012); and *G. hederacea* is toxic to some species of herbivorous insects as it produces a defensive insecticide protein (Hutchings & Price, 1999; Van Damme, 2008). It is also important to note that the negative effect we found did not persist for *B. pascuorum* once our values for *Lamium album* were replaced with higher values found in the literature. Our methods for extracting and measuring protein content are crude, and results can be variable, so should be treated with caution. More detailed investigations are needed in which the full range of compounds present in pollen are quantified if we are to fully understand how bees choose which flowers to visit when collecting pollen.

Most studies do not measure both nectar and pollen rewards in relation to insect visit frequency, and in studies that do, conflicting results have been found. Konzmann and Lunau (2014) found that, in bumblebees, nectar rewards appear more important than pollen quality, whereas Somme *et al.* (2015) found when pollen loads are analysed in conjunction with nectar from forage plants, both nectar and pollen quality appear important. In this study, while pollen and nectar are not negatively correlated, total nectar production appears to influence the

visitation of insects to a greater extent than pollen. This could mean that flower visiting insects are more concerned with the quality of nectar, with pollen as a secondary consideration.

Insect visitation to test plant species appears to be species-specific, which can go further to explain our results. *B. pascuorum* mainly visited *L. album* and *S. officinale*, most likely due to its longer tongue allowing access to their deeper corollas. Contrary to other visiting species, we detected a negative effect of total sugar on *B. pascuorum* visitation. This may be because *B. pascuorum* had little competition from other insect foragers for *L. album* (only 16 other individuals recorded foraging on this species, which produced comparatively low amounts of sugar), or because this species was at the start of its life cycle when this study's sampling was undertaken and newly emerged queens were focusing foraging efforts on pollen collection. *Bombus pratorum*, which as a short-tongued bee may have exhibited restricted foraging choices, tending to visit flowers with greater total sugar. Although this species has a short tongue, it is a secondary nectar robber; 93% of recorded visits to *S. officinale* were via robbing (behaviour previously reported in Goulson *et al.*, 1998b). *Apis mellifera*, although also a short tongued species, mainly visited *R. fruticosus* and was not recorded robbing in this study, though it has been recorded acting as a secondary robber elsewhere (Darwin, 1872). Differences between foraging behaviour of bumblebees and honeybees suggest bumblebees (*B. terrestris* and *B. pascuorum*) show less fidelity when collecting pollen than honeybees, which have a highly flower-constant strategy (Leonhardt & Blüthgen, 2012).

Promoting and developing resources for pollinating insects is predominantly conducted through agri-environmental schemes promoting flower-rich field edges (Carvell *et al.*, 2007), or through targeted planting in urban spaces or private gardens (Hanley *et al.*, 2014). Our understanding of the way in which pollinating insects respond to differences in the quality of resources offered by managed planting is limited. Our results suggest more consideration should be given when selecting plants for conservation management efforts, notably in terms of differing insect species requirements for pollen and nectar quality. Differences in flower selection between pollinator species may relate to the variation of life histories and may reduce competition for resources. Nectar resource quality appears to be the main driver of flower selection by most insect foragers in this study but, importantly, the benefits of greater resource quality in plants are dependent on local floral abundance. One practical conclusion to be drawn from this is that bees may benefit more from plantings of flowers (be they in farmland, parks or gardens) where species are presented in large clumps rather than in heterogeneous mixtures. More broadly, it is clear that there is still much that we do not understand about the role of sugars, proteins and other compounds in nectar and pollen in determining the foraging preferences of pollinators under field conditions.

Acknowledgements

We thank Rebecca Crowe and Charles Linton for help with observation surveys, and the Biotechnology and Biological Sciences Research Council (BBSRC) (BB/J014753/2) for funding this project. We also thank Mick Hanley and three anonymous reviewers for their comments on the original manuscript, which helped improve it considerably.

Supporting Information

Additional Supporting Information may be found in the online version of this article under the DOI reference: doi: 10.1111/icad.12197:

Figure S1. Boxplots illustrating total visits of all insects, all bumblebees, *Bombus pascuorum*, *B. pratorum* and *Apis mellifera* per 10 minute survey from 112 surveys undertaken to measure the influence of nectar and pollen quality on flower visiting insect foraging choice in Sussex, UK.

Table S1. Family, full species name, biological form, and inflorescence type of *Crataegus monogyna*, *Rubus fruticosus*, *Glechoma hederacea*, *Lamium album*, *Ranunculus repens* and *Symphytum officinale*, which were used as test plant species from which to measure nectar and pollen quality and quantity in relation to flower visiting insect foraging choices.

Table S2. The name, Latitude, Longitude and description of sites used for sampling insect foraging visits to *Crataegus monogyna*, *Rubus fruticosus*, *Glechoma hederacea*, *Lamium album*, *Ranunculus repens* and *Symphytum officinale* in Sussex, UK.

Table S3. The number of insects recorded visiting test plant species from 112 observation surveys lasting 10 minutes each conducted in May and June 2014.

Table S4. Linear Mixed Effect (LME) model test results between insect visitation (response) and floral abundance and test species (fixed factors), with sample site as a random factor. Displayed are parameter estimates \pm standard error (SE), *z*-value and *P*-value for each explanatory variable in the final model after simplification, and model AIC and log-likelihood tests.

Table S5. Generalised Linear Mixed Models (GLMM) testing the effects of floral abundance, total nectar and pollen quality on visitation of insect foragers from seven sites in Sussex, UK. These models are the same as Table 2 in the manuscript, apart from the protein value for *Lamium album* changed to 22.8% as reported in Roulston *et al.* (2000) to make sure the low value for this species did not influence model outputs. Only models for *Apis mellifera* and *Bombus pascuorum* are shown as these were the only models with Total Protein left in after model reduction and the only models affected by the change in protein content values. Displayed are parameter estimates \pm standard error, *z*-value and *P*-value for each explanatory variable in the final model after

simplification, and model AIC and log-likelihood tests. *P*-values are in bold if <0.05 significance.

References

- Alaux, C., Ducloz, F., Crauser, D. & Le Conte, Y. (2010) Diet effects on honeybee immunocompetence. *Biology Letters*, **6**, 562–565.
- Arenas, A. & Farina, W.M. (2012) Learned olfactory cues affect pollen-foraging preferences in honeybees, *Apis mellifera*. *Animal Behaviour*, **83**, 1023–1033.
- Basualdo, A.M., Barragán, S., Vanagas, L., García, C., Solana, H., Rodríguez, E. & Bedascarrasbure, E. (2012) Conversion of high and low pollen protein diets into protein in worker honey bees (Hymenoptera: Apidae). *Economic Entomology*, **106**, 1553–1558.
- Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, **67**, 1–48.
- Boggs, C.L. & Ross, C.L. (1993) The effect of adult food limitation on life history traits in *Speyeria mormonia* (Lepidoptera: Nymphalidae). *Ecology*, **74**, 433–441.
- Bolten, A.B., Feinsinger, P., Baker, H.G. & Baker, I. (1979) On the calculation of sugar concentration in flower nectar. *Oecologia*, **41**, 301–304.
- Boppré, M., Colegate, S.M. & Edgar, J.A. (2005) Pyrrolizidine alkaloids of *Echium vulgare* honey found in pure pollen. *Journal of Agricultural and Food Chemistry*, **53**, 594–600.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, **72**, 248–254.
- Brereton, T.M., Roy, D.B., Middlebrook, I., Botham, M. & Warren, M. (2011) The development of butterfly indicators in the United Kingdom and assessments in 2010. *Journal of Insect Conservation*, **15**, 139–151.
- Carvell, C., Meek, W.R., Pywell, R.F., Goulson, D. & Nowakowski, M. (2007) Comparing the efficacy of agri-environment schemes to enhance bumblebee abundance and diversity on arable field margins. *Journal of Applied Ecology*, **44**, 29–40.
- Cook, S.M., Awmack, C.S. & Williams, I.W. (2003) Are honey bees' foraging preferences affected by pollen amino acid composition? *Ecological Entomology*, **28**, 622–627.
- Cox-Foster, D.L., Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.-L., Briese, T., Hornig, M., Geiser, D.M., Martinson, V., vanEngelsdorp, D., Kalkstein, A.L., Drysdale, A., Hui, J., Zhai, J., Cui, L., Hutchison, S.K., Simons, J.F., Egholm, M., Pettis, J.S. & Lipkin, W.I. (2007) A metagenomic survey of microbes in honey bee colony collapse disorder. *Science*, **318**, 283–287.
- Darwin, C. (1872) *The Effects of Cross and Self Fertilisation in the Vegetable Kingdom*. Murray, London, UK.
- Di Pasquale, G., Salignon, M., Le Conte, Y., Belzunces, L.P., Decourtye, A., Kretzschmar, A., Suchail, S., Brunet, J. & Alaux, C. (2013) Influence of pollen nutrition on honey bee health: do pollen quality and diversity matter? *PLoS ONE*, **8**, e72016.
- Dötterl, S. & Vereecken, N.J. (2010) The chemical ecology and evolution of bee–flower interactions: a review and perspectives. *Canadian Journal of Zoology*, **88**, 668–697.
- Geiger, F., Bengtsson, J., Berendse, F., Weisser, W.W., Emmerson, M., Morales, M.B., Ceryngier, P., Liira, J., Tschamtké, T., Winqvist, C., Eggers, S., Bommarco, R., Pärt, T., Bretagnolle, V., Plantegenest, M., Clement, L.W., Dennis, C., Palmer, C., Oñate, J.J., Guerrero, I., Hawro, V., Aavik, T., Thies, C., Flohre, A., Hänke, S., Fischer, C., Goedhart, P.W. & Inchausti, P. (2010) Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology*, **11**, 97–105.
- Génissel, A., Aupinel, P., Bressac, C., Tasei, J.-N. & Chevrier, C. (2002) Influence of pollen origin on performance of *Bombus terrestris* micro-colonies. *Entomologia Experimentalis et Applicata*, **104**, 329–336.
- Gosselin, M., Michez, D., Vanderplanck, M., Roelants, D., Glauser, G. & Rasmont, P. (2013) Does *Aconitum septentrionale* chemically protect floral rewards to the advantage of specialist bumblebees? *Ecological Entomology*, **38**, 400–407.
- Goulson, D., Hanley, M.E., Darvill, B., Ellis, J.S. & Knight, M.E. (2005) Causes of rarity in bumblebees. *Biological Conservation*, **122**, 1–8.
- Goulson, D., Hawson, S.A. & Stout, J.C. (1998a) Foraging bumblebees avoid flowers already visited by conspecifics or by other bumblebee species. *Animal Behaviour*, **55**, 199–206.
- Goulson, D., Lye, G. & Darvill, B. (2008) Decline and conservation of bumble bees. *Annual Review of Entomology*, **53**, 191–208.
- Goulson, D., Nicholls, B., Botías, C. & Rotheray, E.L. (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, **347**, 6229.
- Goulson, D., Stout, J.C., Hawson, S.A. & Allen, J.A. (1998b) Floral display size in comfrey, *Symphytum officinale* L. (Boraginaceae): relationships with visitation by three bumblebee species and subsequent seed set. *Oecologia*, **113**, 502–508.
- Hanley, M.E., Awbi, A.J. & Franco, M. (2014) Going native? Flower use by bumblebees in English urban gardens. *Annals of Botany*, **113**, 799–806.
- Hanley, M.E., Franco, M., Pichon, S., Darvill, B. & Goulson, D. (2008) Breeding system, pollinator choice and variation in pollen quality in British herbaceous plants. *Functional Ecology*, **22**, 592–598.
- Hanley, M.E. & Wilkins, J.P. (2015) On the verge? Preferential use of road-facing hedgerow margins by bumblebees in agroecosystems. *Journal of Insect Conservation*, **19**, 67–74.
- Haughton, A.J., Champion, G.T., Hawes, C., Heard, M.S., Brooks, D.R., Bohan, D.A., Clark, S.J., Dewar, A.M., Firbank, L.G., Osborne, J.L., Perry, J.N., Rothery, P., Roy, D.B., Scott, R.J., Woivod, I.P., Birchall, C., Skellern, M.P., Walker, J.H., Baker, P., Browne, E.L., Dewar, A.J.G., Garner, B.H., Haylock, L.A., Horne, S.L., Mason, N.S., Sands, R.J.N. & Walker, M.J. (2003) Invertebrate responses to the management of genetically modified herbicide-tolerant and conventional spring crops. II. Within-field epigeal and aerial arthropods. *Philosophical transactions of the Royal Society of London Series B, Biological Sciences*, **358**, 1847–1862.
- Heil, M. (2011) Nectar: generation, regulation and ecological functions. *Trends in Plant Science*, **16**, 191–200.
- Heinrich, B. (1979a) Resource heterogeneity and patterns of movement in foraging bumblebees. *Oecologia*, **40**, 235–245.
- Heinrich, B. (1979b) “Majoring” and “Minoring” by foraging bumblebees, *Bombus vagans*: an experimental analysis. *Ecology*, **60**, 245–255.
- Hendriksma, H.P., Oxman, K.L. & Shafir, S. (2014) Amino acid and carbohydrate tradeoffs by honey bee nectar foragers and their implications for plant-pollinator interactions. *Journal of Insect Physiology*, **69**, 56–64.

- Howell, A.D. & Alarcón, R. (2007) *Osmia* bees (Hymenoptera: Megachilidae) can detect nectar-rewarding flowers using olfactory cues. *Animal Behaviour*, **74**, 199–205.
- Hutchings, M.J. & Price, E.A. (1999) *Glechoma hederacea* L. (*Nepeta glechoma* Benth., *N. hederacea* (L.) Trev.). *Ecology*, **87**, 347–364.
- Jurgens, A. & Dötterl, S. (2004) Chemical composition of anther volatiles in Ranunculaceae: genera-specific profiles in *Anemone*, *Aquilegia*, *Caltha*, *Pulsatilla*, *Ranunculus*, and *Trollius* species. *American Journal of Botany*, **91**, 1969–1980.
- Kessler, A. & Halitschke, R. (2009) Testing the potential for conflicting selection on floral chemical traits by pollinators and herbivores: predictions and case study. *Functional Ecology*, **23**, 901–912.
- Kim, Y. & Smith, B. (2000) Effect of an amino acid on feeding preferences and learning behavior in the honey bee, *Apis mellifera*. *Journal of Insect Physiology*, **46**, 793–801.
- Kipling, R.P. & Warren, J. (2013) How generalists coexist: the role of floral phenotype and spatial factors in the pollination systems of two *Ranunculus* species. *Journal of Plant Ecology*, **7**, 1–10.
- Kitaoka, T.K. & Nieh, J.C. (2009) Bumble bee pollen foraging regulation: role of pollen quality, storage levels, and odor. *Behavioral Ecology Sociobiology*, **63**, 625.
- Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S., Kremen, C. & Tscharntke, T. (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B-Biological Sciences*, **274**, 303–313.
- Konzmann, S. & Lunau, K. (2014) Divergent rules for pollen and nectar foraging bumblebees - a laboratory study with artificial flowers offering diluted nectar substitute and pollen surrogate. *PLoS ONE*, **9**, e91900.
- Kremen, C., Williams, N.N. & Thorp, R.W. (2002) Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences of the United States of America*, **99**, 16812–16816.
- Lack, A.J. (1982) Competition for pollinations in the ecology of *Centaurea scabiosa* L. and *Centaurea nigra* L. III. Insect visits and the number of successful pollinations. *New Phytologist*, **91**, 321–339.
- Leonhardt, S.D. & Blüthgen, N. (2012) The same, but different: pollen foraging in honeybee and bumblebee colonies. *Apidologie*, **43**, 449–464.
- Levin, M.D. & Bohart, G.E. (1955) Selection of pollens by honey bees. *American Bee Journal*, **95**, 392–393.
- Maćukanović-Jocić, M., Duletić-Laušević, S. & Jicić, G. (2004) Nectar production in three melliferous species of *Lamiaceae* in natural and experimental conditions. *Acta Veterinaria*, **54**, 475–487.
- Maćukanović-Jocić, M. & Đurđević, L. (2005) Influence of microclimatic conditions on nectar exudation in *Glechoma hirsuta* W.K. *Archives of Biological Sciences*, **57**, 119–126.
- Naimi, B. (2013) *Usdm: Uncertainty Analysis for Species Distribution Models (Version 1.1–12)*. <<http://cran.r-project.org/package=usdm>> 1st October 2015.
- Nicolson, S.W. (2011) Bee Food: The chemistry and nutritional value of nectar, pollen and mixtures of the two. *African Zoology*, **46**, 197–204.
- Öckinger, E. & Smith, H.G. (2007) Semi-natural grasslands as population sources for pollinating insects in agricultural landscapes. *Journal of Applied Ecology*, **44**, 50–59.
- O'Hara, R.B. & Kotze, D.J. (2010) Do not log-transform count data. *Methods in Ecology and Evolution*, **1**, 118–122.
- Ollerton, J., Erenler, H., Edwards, M. & Crockett, R. (2014) Extinctions of aculeate pollinators in Britain and the role of large-scale agricultural changes. *Science*, **346**, 1360–1362.
- Pernal, S.F. & Currie, R.W. (2001) The influence of pollen quality on foraging behavior in honeybees (*Apis mellifera* L.). *Behavioral Ecology Sociobiology*, **51**, 53–68.
- Pinheiro, L.A., Torres, L.M., Raimundo, J. & Santos, S.A.P. (2015) Effects of pollen, sugars and honeydew on lifespan and nutrient levels of *Episyrphus balteatus*. *BioControl*, **60**, 47–57.
- Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O. & Kunin, W.E. (2010) Global pollinator declines: trends, impacts and drivers. *Trends in Ecology and Evolution*, **25**, 345–353.
- Pyke, G.H. (1978) Optimal foraging: movement patterns of bumblebees between inflorescences. *Theoretical Population Biology*, **13**, 72–98.
- R Development Core Team (2014) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <<http://www.R-project.org>> 1st October 2015.
- Robertson, A.W., Mountjoy, C., Faulkner, B.E., Roberts, V.M. & Macnair, M.R. (1999) Bumble bee selection of *Mimulus guttatus* flowers: the effects of pollen quality and reward depletion. *Ecology*, **80**, 2594–2606.
- Robinson, R.A. & Sutherland, W.J. (2002) Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology*, **39**, 157–176.
- Roulston, T.H. & Cane, J.H. (2000) Pollen nutritional content and digestibility for animals. *Plant Systematics and Evolution*, **222**, 187–209.
- Roulston, T.H. & Cane, J.H. (2002) The effect of pollen protein concentration on body size in the sweat bee *Lasioglossum zephyrum* (Hymenoptera: Apiformes). *Evolutionary Ecology*, **16**, 49–65.
- Roulston, T.H., Cane, J.H. & Buchmann, S.L. (2000) What governs protein content of pollen: pollinator preferences, pollen-pistil interactions, or phylogeny? *Ecological Monographs*, **70**, 617–643.
- RStudio (2012) *RStudio: Integrated Development Environment for R (Version 0.98.1091)*. <<http://www.rstudio.org/>> 11th June 2015.
- Schmidt, J.O. (1982) Pollen foraging preferences of honey bees. *The Southwestern Entomologist*, **7**, 255–259.
- Sedivy, C., Piskorski, R., Müller, A. & Dorn, S. (2012) Too low to kill: concentration of the secondary metabolite ranunculin in buttercup pollen does not affect bee larval survival. *Journal of Chemical Ecology*, **38**, 996–1002.
- Silva, E.M. & Dean, B.B. (2004) Patterns of floral nectar production of Onion (*Allium cepa* L.) and the effects of environmental conditions. *Journal of the American Society for Horticultural Science*, **129**, 299–302.
- Somme, L., Vanderplanck, M., Michez, D., Lombaerde, I., Moerman, R., Wathélet, B., Wattiez, R., Lognay, G. & Jacquemart, A.-L. (2015) Pollen and nectar quality drive the major and minor floral choices of bumble bees. *Apidologie*, **46**, 92–106.
- Stone, G.N. (1993) Endothermy in the solitary bee *Anthophora plumipes*: independent measures of thermoregulatory ability, costs of warm-up and the role of body size. *Journal of Experimental Biology*, **174**, 299–320.

- Strohm, E. & Linsenmair, K.E. (1997) Female size affects provisioning and sex allocation in a digger wasp. *Animal Behaviour*, **54**, 23–34.
- Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.S. (2008) Global change and species interactions in terrestrial ecosystems. *Ecology Letters*, **11**, 1351–1363.
- Van Damme, E.J. (2008) Plant lectins as part of the plant defense system against insects. *Induced plant resistance to herbivory*, pp. 285–307. (ed. by A. Schaller), Springer, Netherlands.
- Waser, N.M., Chittka, L., Price, M.V., Williams, N.M. & Ollerton, J. (1996) Generalization in pollination systems, and why it matters. *Ecology*, **77**, 1043–1060.
- Wickham, H. (2009) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York City, New York.
- Wille, H., Wille, M., Kilchenmann, V., Imdorf, A. & Bühlmann, G. (1985) Pollenernte und Massenwechsel von drei *Apis mellifera*-Völkern auf demselben Bienenstand in zwei aufeinanderfolgenden Jahren. *Revue Suisse de Zoologie*, **92**, 897–914. In: Praz, C., Müller, A. & Dorn, S. (2008) Specialized bees fail to develop on non-host pollen: Do plants chemically protect their pollen? *Ecology*, **89**, 795–804.
- Wolff, D. (2006) Nectar sugar composition and volumes of 47 species of Gentianales from a southern Ecuadorian montane forest. *Annals of Botany*, **97**, 767–777.
- Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010) A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, **1**, 3–14.

Accepted 12 August 2016

Editor: Karsten Schonrogge
Associate editor: Matt Heard